



# FC Professional Insect Control™ Ant Killer Bait Gel

## FOR USE IN COMMERCIAL, INDUSTRIAL AND RESIDENTIAL AREAS

### ACTIVE INGREDIENT:

Fipronil† .....0.001%

OTHER INGREDIENTS: .....99.999%

†CAS NO. 120068-37-3

Contains: 3 - 60 gm plastic syringes

### KEEP OUT OF REACH OF CHILDREN

### CAUTION: SEE CAUTIONS BELOW

Use only in areas not easily accessible to children and pets.

Product may stain textiles and clothing.

This product provides FAST CONTROL of ants in 3-5 days. MAXFORCE® FC PROFESSIONAL INSECT CONTROL™ ANT KILLER BAIT GEL is formulated to effectively control ants. That's because this product contains Fipronil, a fast-acting active ingredient. Its speed is still slow enough, however, for foraging ants to return to the colony and feed the bait to larvae and queen(s) so that the entire colony is killed through the Domino Effect.™

**This product is formulated for effective ant control.** This product contains a unique blend of inert ingredients specifically designed for ants. The high moisture content of this gel enhances the bait's attractiveness to target ants. The bait's attractiveness is greatest within two weeks of application. This product is a bait product, formulated to take advantage of ant trophallaxis. Through trophallaxis, worker ants eat the bait while foraging, then feed it to the rest of the colony. The larvae digest the solid bait and produce a liquid containing the Fipronil that is fed to the rest of the colony and queen(s).

**This product controls a broad spectrum of ants.** This product's unique blend of inert ingredients is attractive to a broad spectrum of ants. Controls ants including Argentine, Pharaoh, Pavement, Ghost, Honey, Acrobat, Odorous House, Thief, Big-headed, Cornfield, and Field.

[When exposed to higher temperatures (above 80°F) for extended periods of time, the gel will change color and lose effectiveness. If the gel outside the tube is darker than the color bar below, do not apply.]

### PRECAUTIONARY STATEMENTS: HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** May be harmful if swallowed. Avoid contact with skin and clothing. Keep exposed gel away from open foods and food contact surfaces. Wash hands thoroughly after handling.

**FIRST AID:** If swallowed, drink two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. In case of emergency, call day or night, 1-800-334-7577.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a cool, dry area out of reach of children.

**DISPOSAL:** Wrap used prefilled reservoir in several layers of newspaper and discard in trash.

For questions or comments about MAXFORCE, call toll-free 1-800-331-2867.

### DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application rate is dependent on level of infestation and species to be controlled. Bait should be applied as spots.

**Applications in Food/Feed Handling Establishments. Application within food/feed areas of food/feed handling establishments is limited to crack and crevice treatment only.** Apply with syringe applicator or bait injector directly into cracks and crevices. Place applicator tip into cracks and crevices and where ants may travel. Apply gel (up to 1 gram) per spot. Apply bait in small amounts between different elements of construction, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, cabinets, sinks, stoves and other places where ants may enter the establishment. During follow-up visits inspect bait placements and reapply when necessary.

Do not apply bait to areas where food/feed, food utensils or food processing surfaces may become contaminated. Do not apply bait in cracks and crevices of areas that are routinely washed, tops of tables, food/feed preparation and prepared food/feed holding surfaces as bait may be removed by washing. Do not apply bait to surfaces where the temperature exceeds 120°F as bait may run. Examples include, but are not limited to, portions of stoves, ovens, grills, fume hoods, heat lamps, coffee urns, steam tables, toasters, fryers, dishwashers and hot water pipes. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove it and wash exposed surface.

Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered a food area when food is exposed and facility is in operation. **Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted.**

1. To apply gel: Remove cap from tip, touch tip to surface and depress plunger. Recap after use.

2. Apply gel in spots in cracks and crevices. For best results, make -or- place many spots of dabs of gel where you see ants foraging and entering the structure.

3. Do not apply the gel to areas which have been recently sprayed with insecticide or contaminated with foggers, and do not spray insecticide over gel as this may cause the bait to become repellent. Do not apply gel to areas that are routinely washed, such as counter tops, as gel will be removed or diluted and spread onto the surface. If you have recently sprayed, remove the dead ants and thoroughly clean the area before applying the gel.

4. Place the bait as follows: Gel can be placed in cracks, crevices, corners and any out of the way place, such as under and behind kitchen appliances, along baseboards, under sinks, in pantry, around pipes and water heater, window and door frames, and in the garage, attic, or crawl space.

**For indoor use:** Visually inspect for ants and place gel in areas where you see ants. Likely locations for ants and gel placements are areas where ants enter the building structure and areas with potential water damage including: wall voids, cracks and crevices in kitchen, bathroom, under sink, window sills, screen vents, garage, attic, and crawl spaces. Do not place bait in locations that are routinely washed, as bait will be removed by washing. Do not treat food preparation surfaces. Do not use this product in or on electrical equipment where a possibility of shock hazard exists.

**For outdoor use:** Place gel outdoors into, on or adjacent to structures where you see ants, next to ant trails and suspected areas of ant activity. Likely locations for ants and gel placements are points of entry into structure and areas with potential water damage including: eaves, soffits, weep holes, where wires and pipes come into the building structure, trees, stumps, roots, wood decks, fences, planter boxes, woodpiles, landscaping, and railroad ties. Extreme heat or cold, direct sunlight and water may reduce the residual effectiveness. Apply Gel outdoors into MAXFORCE Refillable Bait Stations or other devices designed to hold and protect bait outdoors. Close Refillable MAXFORCE Bait Station securely after application.

Inspect placements and reapply as needed; The gel will remain attractive to ants for up to two weeks. Inspect gel placements and reapply as needed. reapply in the same area if bait is gone or in a different location if no ant activity is evident.

Pest population reduction will be apparent within a week.

Maxforce is a registered trademark of Bayer AG.  
FC Professional Insect Control is a trademark of Bayer AG.

Contains 3 plastic syringes  
NET WEIGHT 2.12 oz (60g), TOTAL NET WEIGHT 6.3 oz. (108 g)



### FC Professional Insect Control™ Ant Killer Bait Gel

Bayer Environmental Science  
A Business Group of Bayer CropScience LP  
95 Chestnut Ridge Road, Montvale, NJ 07645

EPA Reg. No. 432-1264  
EPA Est. No. 5813-MS-001  
3/03